



**If you experience any of the symptoms of moderately to severely active RA, make sure you speak to a rheumatologist as soon as possible. Remember, only a qualified physician can make a diagnosis and help you make an appropriate treatment choice.**

As you speak to your doctor about how RA may be affecting you, use the following questions to guide your discussion. When the discussion turns to your treatment options, ask your doctor if REMICADE® is right for you.

REMICADE®, used in combination with methotrexate, can reduce signs and symptoms, help stop further joint damage, and improve physical function in patients with moderately to severely active rheumatoid arthritis. **Please read the Important Safety Information for REMICADE® below.**

**Discuss these questions and answers with your rheumatologist:**

1. Are you taking any medications on a regular basis for RA?\* (Please check all that apply)

- |  |   |
|--|---|
| <input type="checkbox"/> Arava® (leflunomide)            | <input type="checkbox"/> Humira® (adalimumab)                           |
| <input type="checkbox"/> Neoral® (cyclosporine)          | <input type="checkbox"/> Actemra® (tocilizumab)                         |
| <input type="checkbox"/> Plaquenil® (hydroxychloroquine) | <input type="checkbox"/> Celebrex® (celecoxib)                          |
| <input type="checkbox"/> Methotrexate                    | <input type="checkbox"/> Steroids (eg, prednisone)                      |
| <input type="checkbox"/> Cimzia® (certolizumab pegol)    | <input type="checkbox"/> Advil® (ibuprofen) or Aleve® (naproxen sodium) |
| <input type="checkbox"/> Enbrel® (etanercept)            | <input type="checkbox"/> Other  |

\*Indicated trademarks are registered trademarks of their respective owners.

2. Are you getting the relief you want from your medication?

- No     Rarely     Occasionally     Frequently

3. Even with your current medication, are you still experiencing painful, stiff, and swollen joints?

- No     Rarely     Occasionally     Frequently

4. Has your joint pain increased in severity since you began your current treatment?

- Yes     No

5. Has the number of joints affected by arthritis pain increased since you began your current treatment?

- Yes     No

6. Describe your level of difficulty in accomplishing the following tasks:

- |                           |   |   |
|---------------------------|---|---|
| Dressing yourself         | <input type="checkbox"/> Without any difficulty | <input type="checkbox"/> With some difficulty |
|                           | <input type="checkbox"/> With much difficulty   | <input type="checkbox"/> Unable to do         |
| Gripping a cup            | <input type="checkbox"/> Without any difficulty | <input type="checkbox"/> With some difficulty |
|                           | <input type="checkbox"/> With much difficulty   | <input type="checkbox"/> Unable to do         |
| Shampooing your hair      | <input type="checkbox"/> Without any difficulty | <input type="checkbox"/> With some difficulty |
|                           | <input type="checkbox"/> With much difficulty   | <input type="checkbox"/> Unable to do         |
| Climbing a few stairs     | <input type="checkbox"/> Without any difficulty | <input type="checkbox"/> With some difficulty |
|                           | <input type="checkbox"/> With much difficulty   | <input type="checkbox"/> Unable to do         |
| Getting in and out of bed | <input type="checkbox"/> Without any difficulty | <input type="checkbox"/> With some difficulty |
|                           | <input type="checkbox"/> With much difficulty   | <input type="checkbox"/> Unable to do         |

7. In spite of your current medication, are you still frustrated by not being able to do daily activities?

- No
- Rarely
- Occasionally
- Frequently

8. If I'm feeling better and not having as much pain or stiffness in my joints, does that mean that RA is not damaging my joints?

9. Can REMICADE® help stop further joint damage from RA?

### Have some questions of your own?

If you have more questions you'd like to ask, use the space below to write them down. It's always best to prepare your questions before your appointment so you don't forget anything. Let your doctor know at the beginning of the appointment that you have some questions so there will be time to answer all of them.

---

---

---

---

---

---

---

---

---

---

## IMPORTANT SAFETY INFORMATION FOR REMICADE® (INFLIXIMAB)

Only your doctor can recommend a course of treatment after checking your health condition. REMICADE® (infliximab) can cause serious side effects such as lowering your ability to fight infections. **Some patients, especially those 65 years and older, have had serious infections caused by viruses, fungi or bacteria that have spread throughout the body, including tuberculosis (TB) and histoplasmosis. Some of these infections have been fatal. Your doctor should monitor you closely for signs and symptoms of TB during treatment with REMICADE®.**

**Unusual cancers have been reported in children and teenage patients taking TNF-blocker medicines. Hepatosplenic T-cell lymphoma, a rare form of fatal lymphoma, has occurred mostly in teenage or young adult males with Crohn's disease or ulcerative colitis who were taking REMICADE® and azathioprine or 6-mercaptopurine.** For children and adults taking TNF blockers, including REMICADE®, the chances of getting lymphoma or other cancers may increase.

You should discuss any concerns about your health and medical care with your doctor.

### What should I tell my doctor before I take REMICADE®?

You should let your doctor know if you have or ever had any of the following:

- Tuberculosis (TB) or have been near someone who has TB. Your doctor will check you for TB with a skin test. If you have latent (inactive) TB, you will begin TB treatment before you start REMICADE®.
- Lived in a region where certain fungal infections like histoplasmosis or coccidioidomycosis are common.
- Infections that keep coming back, have diabetes or an immune system problem.
- Any type of cancer or a risk factor for developing cancer, for example, chronic obstructive pulmonary disease (COPD) or had phototherapy for psoriasis.
- Heart failure or any heart condition. Many people with heart failure should not take REMICADE®.
- Hepatitis B virus (HBV) infection or think you may be a carrier of HBV. Your doctor will test you for HBV.
- Nervous system disorders (like multiple sclerosis or Guillain-Barré syndrome).

Also tell your doctor about any medications you are taking, including vaccines or Kineret (anakinra), Orencia (abatacept) or Actemra (tocilizumab) and if you are pregnant, plan to become pregnant or are nursing. Adults and children should not receive a live vaccine while taking REMICADE®. If you have a baby and you were using REMICADE® during your pregnancy, tell your baby's doctor about your REMICADE® use before the baby receives any vaccine.

### What should I watch for and talk to my doctor about before or while taking REMICADE®?

The following serious (sometimes fatal) side effects have been reported in people taking REMICADE®.

You should tell your doctor right away if you have any of the signs listed below:

- Infections (like TB, blood infections, pneumonia)—fever, tiredness, cough, flu, or warm, red or painful skin or any open sores. REMICADE® can make you more likely to get an infection or make any infection that you have worse.
- Lymphoma, or any other cancers in adults and children.
- Heart failure—new or worsening symptoms, such as shortness of breath, swelling of your ankles or feet, or sudden weight gain.
- Reactivation of HBV—feeling unwell, poor appetite, tiredness, fever, skin rash and/or joint pain.
- Liver injury—jaundice (yellow skin and eyes), dark brown urine, right-sided abdominal pain, fever, or severe tiredness.

*(Continued...)*

## IMPORTANT SAFETY INFORMATION FOR REMICADE® (INFLIXIMAB) (continued)

- Blood disorders—fever that doesn't go away, bruising, bleeding or severe paleness.
- Nervous system disorders—numbness, weakness, tingling, changes in your vision or seizures.
- Allergic reactions during or after the infusion—hives, difficulty breathing, chest pain, high or low blood pressure, swelling of face and hands, and fever or chills.
- Lupus-like syndrome—chest discomfort or pain that does not go away, shortness of breath, joint pain, rash on the cheeks or arms that gets worse in the sun. The more common side effects with REMICADE® are respiratory infections (that may include sinus infections and sore throat), headache, rash, coughing and stomach pain.
- Psoriasis—new or worsening psoriasis such as red scaly patches or raised bumps on the skin that are filled with pus.

Click here to read the [Medication Guide](#) for REMICADE® below and discuss it with your doctor.  
Click here for full [Prescribing Information](#).

You are encouraged to report negative side effects of prescription drugs to the FDA.  
Visit [www.fda.gov/medwatch](http://www.fda.gov/medwatch), or call 1-800-FDA-1088.

25R11048