

WELCOME TO REMICADE® (infliximab)

SELECTED IMPORTANT SAFETY INFORMATION

REMICADE® can lower your ability to fight infections. Serious and sometimes fatal events can occur. There have been reports of serious infections including tuberculosis (TB) and infections caused by bacteria, fungi, or viruses that have spread throughout the body. Lymphoma, including a fatal kind called hepatosplenic T-cell lymphoma, and other cancers have been reported in children and adults taking REMICADE®. Some people with heart failure should not take REMICADE®. Other serious side effects reported include skin cancer, hepatitis B, liver injury, blood problems, allergic reactions, nervous system problems, or lupus-like syndrome. To learn more about these and other risks, please read the Important Safety Information and the Medication Guide and discuss any questions you have with your doctor.



The person depicted is a model used for illustrative purposes only.

WELCOME TO A TREATMENT THAT OFFERS PROVEN SYMPTOM CONTROL WELCOME TO REMICADE®

SELECTED IMPORTANT SAFETY INFORMATION

REMICADE® can lower your ability to fight infections. Serious and sometimes fatal events can occur. There have been reports of serious infections including tuberculosis (TB) and infections caused by bacteria, fungi, or viruses that have spread throughout the body. Lymphoma, including a fatal kind called hepatosplenic T-cell lymphoma, and other cancers have been reported in children and adults taking REMICADE®. Some people with heart failure should not take REMICADE®. Other serious side effects reported include skin cancer, hepatitis B, liver injury, blood problems, allergic reactions, nervous system problems, or lupus-like syndrome. To learn more about these and other risks, please read the [Important Safety Information](#) and the [Medication Guide](#) and discuss any questions you have with your doctor.

Janssen CarePath Savings Program for REMICADE®

GETTING STARTED

You and your doctor will fill out the Janssen CarePath Savings Program enrollment form. After you enroll, a Janssen CarePath Savings Program card will be sent to you to pay for your medication out-of-pocket cost.

GETTING YOUR REBATE

- STEP 1** → Receive your REMICADE® (infliximab) intravenous (IV) infusion at your doctor's office or infusion center and pay your co-pay to the office.
- STEP 2** → The doctor's office or infusion center will submit a claim for your infusion to your insurance company.
- STEP 3** → Both you and your doctor's office or infusion center will receive an Explanation of Benefits (EOB) statement from your insurance company. Fax your EOB to Janssen CarePath Savings Program at **1-877-234-3048**.
- STEP 4** → Once the Janssen CarePath Savings Program care coordinator receives your EOB, they will review the information and apply the \$5 rebate to your Janssen CarePath Savings Program card.
- STEP 5** → Your Janssen CarePath Savings Program card is loaded and ready for use at your doctor's office or infusion center to pay for your medication out-of-pocket cost.

Eligible patients enrolled in the Janssen CarePath Savings Program

PAY JUST \$5 PER INFUSION**

Visit Remicade.JanssenCarePathSavings.com for more information and to download forms.

*This program is only available to individuals using private or commercial health insurance to cover a portion of medication costs. This program is not available to individuals who use any state or federal government subsidized healthcare program to cover a portion of medication costs, such as Medicare, Medicaid, TRICARE, Department of Defense, or Veterans Administration. Patients confirm that they will not seek reimbursement from any of these programs or from pharmaceutical patient assistance foundations and accounts such as a Flexible Spending Account (FSA), Healthcare Savings Account (HSA) or Health Reimbursement Account (HRA).

†Eligible patients pay \$5 per infusion, with up to \$20,000 maximum benefit each calendar year. For Massachusetts residents only, this offer is subject to change per state legislation.

‡Janssen CarePath Savings Program rebates are determined by medication cost only. Rebate amounts are not determined by costs associated with administration of the IV infusion.



MAPPING OUT THE IV INFUSION PROCESS

Your doctor's office or infusion center will schedule your appointments. If you need to reschedule an appointment, call your doctor's office or infusion provider.

BEFORE YOU ARRIVE



DRINK UP!

It's a good idea to show up to your IV infusion well hydrated—drinking plenty of water might also help with your symptoms.



PACK UP!

REMICADE® (infliximab) is administered over a period of 2 hours. Bring a book, laptop, tablet, or other digital device...or just a pillow and take a nap to pass the time.



DRESS UP!

It's important to dress for the occasion. Make comfort a priority: wear loose-fitting clothing and layers in case you get too warm or too cold.



Watch a video and learn more about the IV infusion process at [REMICADE.com](https://www.remicade.com)

DURING YOUR IV INFUSION



When you arrive, your healthcare professional will check your vital signs, and ask you about certain health issues, including:

- Any changes in your medical history
- New medications you may be taking
- Recent flu shots or other vaccines
- Any possibility of pregnancy
- Any infection or other symptoms you've noticed
- Heart conditions
- Nervous system disorders



Your hand or arm will be sterilized with rubbing alcohol and an IV needle will be inserted and fastened with tape. Your doctor may decide to give you other medications before starting the REMICADE® infusion to prevent or lessen side effects, some of which may cause drowsiness.



A healthcare professional will check in with you periodically throughout your IV infusion. The IV infusion takes about 2 hours.

AFTER YOUR IV INFUSION

You will be monitored at the doctor's office or infusion center for about 30 minutes just to make sure you are not having a reaction, and then you will be free to leave. REMICADE® does not cause drowsiness so you should be able to continue your day as normal.

REMICADE® isn't right for everyone, and individual results may vary. Talk with your doctor to decide if REMICADE® is right for you.

Please read the [Important Safety Information for REMICADE®](#) and the [Medication Guide](#) and discuss any questions you have with your doctor.



For adult patients with moderately to severely active Crohn's disease or ulcerative colitis who haven't responded well to other therapies

OVER 1 MILLION PATIENTS WITH CROHN'S AND UC HAVE BEEN PRESCRIBED REMICADE® WORLDWIDE*

SELECTED IMPORTANT SAFETY INFORMATION

REMICADE® can lower your ability to fight infections. Serious and sometimes fatal events can occur. There have been reports of serious infections including tuberculosis (TB) and infections caused by bacteria, fungi, or viruses that have spread throughout the body. Lymphoma, including a fatal kind called hepatosplenic T-cell lymphoma, and other cancers have been reported in children and adults taking REMICADE®. Some people with heart failure should not take REMICADE®. Other serious side effects reported include skin cancer, hepatitis B, liver injury, blood problems, allergic reactions, nervous system problems, or lupus-like syndrome. To learn more about these and other risks, please read the Important Safety Information and the Medication Guide and discuss any questions you have with your doctor.

FIND REMICADE® ONLINE!

Want more information about REMICADE® (infliximab)? Maybe you just want to hear from actual patients or learn more about Crohn's disease or UC.

Our Pinterest and YouTube pages have information to help make an informed decision about treatment with REMICADE® in consultation with your doctor.



REMICADE® isn't right for everyone, and individual results may vary. Talk with your doctor to decide if REMICADE® is right for you.

*As of August 2014.



INDICATIONS AND IMPORTANT SAFETY INFORMATION

INDICATIONS

REMICADE® (infliximab) can reduce signs and symptoms and induce and maintain remission in adult patients with moderately to severely active Crohn's disease who haven't responded well to other therapies.

REMICADE® can reduce signs and symptoms, induce and maintain remission, promote intestinal healing, and reduce or stop the need for steroids in adult patients with moderately to severely active ulcerative colitis (UC) who haven't responded well to other therapies.

IMPORTANT SAFETY INFORMATION

Only your doctor can recommend a course of treatment after checking your health condition. REMICADE® can cause serious side effects such as lowering your ability to fight infections. **Some patients, especially those 65 years and older, have had serious infections caused by viruses, fungi or bacteria that have spread throughout the body, including tuberculosis (TB) and histoplasmosis. Some of these infections have been fatal. Your doctor should monitor you closely for signs and symptoms of TB during treatment with REMICADE®.**

Unusual cancers have been reported in children and teenage patients taking TNF-blocker medicines. Hepatosplenic T-cell lymphoma, a rare form of fatal lymphoma, has occurred mostly in teenage or young adult males with Crohn's disease or ulcerative colitis who were taking REMICADE® and azathioprine or 6-mercaptopurine. For children and adults taking TNF blockers, including REMICADE®, the chances of getting lymphoma or other cancers may increase.

You should discuss any concerns about your health and medical care with your doctor.

What should I tell my doctor before I take REMICADE®?

You should let your doctor know if you have or ever had any of the following:

- Tuberculosis (TB) or have been near someone who has TB. Your doctor will check you for TB with a skin test. If you have latent (inactive) TB, you will begin TB treatment before you start REMICADE®.
- Lived in a region where certain fungal infections like histoplasmosis or coccidioidomycosis are common.
- Infections that keep coming back, have diabetes or an immune system problem.
- Any type of cancer or a risk factor for developing cancer, for example, chronic obstructive pulmonary disease (COPD) or had phototherapy for psoriasis.
- Heart failure or any heart condition. Many people with heart failure should not take REMICADE®.
- Hepatitis B virus (HBV) infection or think you may be a carrier of HBV. Your doctor will test you for HBV.
- Nervous system disorders (like multiple sclerosis or Guillain-Barré syndrome).

Also tell your doctor if you:

- Use the medicines Kineret (anakinra), Orencia (abatacept) or Actemra (tocilizumab) or other medicines called biologics used to treat the same problems as REMICADE®.
- Are pregnant, plan to become pregnant, are breast-feeding, or have a baby and were using REMICADE® during your pregnancy. Tell your baby's doctor about your REMICADE® use. If your baby receives a live vaccine within 6 months after birth, your baby may develop infections with serious complications that can lead to death.
- Recently received or are scheduled to receive a vaccine. Adults and children taking REMICADE® should not receive live vaccines or treatment with a weakened bacteria (such as BCG for bladder cancer) while taking REMICADE®.

What should I watch for and talk to my doctor about before or while taking REMICADE®?

The following serious (sometimes fatal) side effects have been reported in people taking REMICADE®.

You should tell your doctor right away if you have any of the signs listed below:

- Infections (like TB, blood infections, pneumonia)—fever, tiredness, cough, flu, or warm, red or painful skin or any open sores. REMICADE® can make you more likely to get an infection or make any infection that you have worse.
- Lymphoma, or any other cancers in adults and children.
- Skin cancer—any changes in or growths on your skin.
- Heart failure—new or worsening symptoms, such as shortness of breath, swelling of your ankles or feet, or sudden weight gain.
- Reactivation of HBV—feeling unwell, poor appetite, tiredness, fever, skin rash and/or joint pain.
- Liver injury—jaundice (yellow skin and eyes), dark brown urine, right-sided abdominal pain, fever, or severe tiredness.
- Blood disorders—fever that doesn't go away, bruising, bleeding or severe paleness.
- Nervous system disorders—numbness, weakness, tingling, changes in your vision or seizures.
- Allergic reactions during or after the infusion—hives, difficulty breathing, chest pain, high or low blood pressure, swelling of face and hands, and fever or chills.
- Lupus-like syndrome—chest discomfort or pain that does not go away, shortness of breath, joint pain, rash on the cheeks or arms that gets worse in the sun.
- Psoriasis—new or worsening psoriasis such as red scaly patches or raised bumps on the skin that are filled with pus.

The more common side effects with REMICADE® are respiratory infections (that may include sinus infections and sore throat), headache, rash, coughing and stomach pain.

Please read the [Medication Guide](#) for REMICADE® and discuss it with your doctor.

You are encouraged to report negative side effects of prescription drugs to the FDA.

Visit www.fda.gov/medwatch, or call 1-800-FDA-1088.



Janssen CarePath Savings Program for REMICADE®

Eligible patients enrolled in the **Janssen CarePath Savings Program**

**PAY JUST \$5
PER INFUSION****

Before the calendar year ends, you will receive information and eligibility requirements for continued benefit in the program.

Visit JanssenCarePath.com/Remicade to see if you qualify, and start saving today!

Janssen CarePath for REMICADE® Support Services

Janssen CarePath provides a variety of services designed to make it easier for you to get the care you need.

Call **877-CarePath** (877-227-3728) Monday-Friday 8 AM - 8 PM ET or visit JanssenCarePath.com/Remicade for more information.

*This program is only available to individuals using private or commercial health insurance to cover a portion of medication costs. This program is not available to individuals who use any state or federal government subsidized healthcare program to cover a portion of medication costs, such as Medicare, Medicaid, TRICARE, Department of Defense, or Veterans Administration. Patients confirm that they will not seek reimbursement from any of these programs or from pharmaceutical patient assistance foundations and accounts such as a Flexible Spending Account (FSA), Healthcare Savings Account (HSA) or Health Reimbursement Account (HRA).

†**Eligible patients pay \$5 per infusion, with up to \$20,000 maximum benefit each calendar year.**
For Massachusetts residents only, this offer is subject to change per state legislation.

‡Janssen CarePath Savings Program rebates are determined by medication cost only. Rebate amounts are not determined by costs associated with administration of the IV infusion.

Please read the Important Safety Information for REMICADE® and the Medication Guide and discuss any questions you have with your doctor.

